

Radiation Field on Superspace[†]

Pedro F. González-Díaz.

Instituto de Matemáticas y Física Fundamental
Consejo Superior de Investigaciones Científicas
Serrano 121, 28006 Madrid (SPAIN)

June 8, 1994

We study the dynamics of multiwormhole configurations within the framework of the Euclidean Polyakov approach to string theory, incorporating a modification to the Hamiltonian which leads to a Planckian probability measure for the Coleman parameters α that allows $\frac{1}{2}\alpha^2$ to be interpreted as the energy of the quanta of a radiation field on superspace whose values might still fix the coupling constants.

[†]To appear in *Geometry of Constrained Dynamical Systems*, ed. J.M. Charap (Cambridge Univ. Press, Cambridge, 1995).

Multiwormhole configurations in Polyakov string theory have been studied by looking at the wormholes as the handles on a Riemann surface of genus g , with g giving the number of handles or wormholes in the configuration [1]. The Green function that describes the effects of such wormholes on first order tachyonic amplitudes was calculated by Lyons and Hawking [2] as a path integral over all space-time coordinates x_μ on the Riemann surface. It was assumed that all the fields have the same values at the points on the two circles which result after cutting the handles in such a way that they become divided in two topologically separated discs. In this case, the points were identified by the projective transformations of the Schottky group on each pair of circles, and the Green function can be written as [1-3]

$$\langle x_\mu(z_1).x_\nu(z_2)\dots \rangle = \int d[x_\mu] x_\mu(z_1)x_\nu(z_2)\dots \prod_r \prod_n \delta(x_n - x'_n) e^{-I}, \quad (1)$$

where I is the Euclidean action, r runs from 1 to g ,

$$x = \sum x_n e^{i\zeta n}, \quad (2)$$

and the delta function ensures that ζ on one circle is identified with ζ' on another. One can express the Green function (1) in terms of the handle quantum state on the circles using the Fourier transform of the delta function for the zero mode, and expanding the delta function for the nonzero modes in terms of the complete set of orthonormal harmonic-oscillator eigenstates which are the solutions of the string analogue of the Wheeler DeWitt equation [2]

$$H_{WDW} \Psi_{nm_n^{(i)}} = \left[-\frac{\partial^2}{\partial x_0^2} + \frac{1}{2} \sum_{n>0,i} \left(-\frac{\partial^2}{\partial (Y_n^{(i)})^2} + n^2 (Y_n^{(i)})^2 \right) \right] \Psi_{nm_n^{(i)}} = 0, \quad (3)$$

with solution ($i = 1, 2$)

$$\Psi_{nm_n^{(i)}} \propto e^{-\frac{1}{2}n(Y_n^{(i)})^2} H_{m_n^{(i)}}(n^{\frac{1}{2}} Y_n^{(i)}) \Psi_K(x_0), \Psi_K(x_0) = e^{iK.x_0}. \quad (4)$$

Eq. (3) is the canonically quantised version of the Hamiltonian constraint derived [2] from the string Euclidean action for the field x on the region of the complex plane outside a disc of radius $r = -\frac{\ln|z|}{t}$ (t is some Euclidean time), K is the momentum of the zero mode $n = 0$, and

$$Y_n^{(1)} = \frac{1}{2}(x_n + x_{-n}), Y_n^{(2)} = \frac{1}{2i}(x_n - x_{-n}).$$

Now, as in the 4-dimensional case [4], one can calculate the effect on Green functions in the fundamental region, by doing a path

integral over all fields x^μ on the complex plane with the boundary conditions on the circles given by a set of values $Y_n^{(i)}$, weighting with the wave function $\Psi_{nm_n}^{(i)}$. Again as in the space-time wormhole case, the effect will be given by a vertex operator located approximately at the center of the circles. One can see that, after integrating over the fields, the resulting path integral contains a factor $\kappa = (K^2 + \sum |n| m_n^{(i)} - 2)^{-1}$. We obtain thus a bi-local effective action [1]

$$- \int d\sigma_1 \int d\sigma_2 \sum \int d^4 K \kappa V_p(\sigma_1) V_p(\sigma_2),$$

where the V_p are the handle vertex operators for the fields on the two circles. One can again convert the bi-local action to a local action, by introducing α parameters, to finally obtain a probability measure with the same general form as for the 4-dimensional case [5]. Nevertheless, since now the α parameters are labelled by momentum K , and also the occupation numbers $m_n^{(i)}$ for modes $n > 0$, these parameters should be interpreted [1,3] as a quantum field on the infinite dimensional superspace of all coordinate field values on a single circle. Such a quantum field would be regarded as an infinite tower of fields on the usual space-time minisuperspace; each stage along this tower is labelled by the quantum number $m_n^{(i)}$ that defines the excited states of the basis set of solutions (4). In light of this interpretation, it was concluded [1,3] that the α 's could not be regarded as a set of coupling constants to be fixed by quantum measurement, as required by the Coleman mechanism to fix the coupling constants [6].

However, it is not quite clear that this conclusion can be maintained because in string theory the Hamiltonian should be modified [7] by the addition of an (infinite) constant, in such a way that just the ground state of the harmonic oscillators, multiplied by the wave function of the zero mode, will obey the Hamiltonian constraint.

We shall follow here a modified approach where this shortcoming is avoided. The idea consists in considering the more general case allowing for handles on the Riemann surface which, under cutting, give rise to pairs of discs that are no longer disconnected to each other [8]. This would ultimately imply partial or total breakdown of the Schottky group invariance under projective transformations between discs [9]. Such handles need not be on shell in the sense that the analogue of the Wheeler DeWitt operator acting on the excited eigenstates $\Psi_{m_n^{(i)}}(Y_n^{(i)})$ is no longer zero, but gives the corresponding harmonic-oscillator eigenvalues $nm_n^{(i)}$ [8,10]. In this case, the δ function for all field modes in the Green function should be replaced [8] by a path integral which has the x_n fixed at the given values at time $t = 0$ on a circle and $t = t_1$ on another, for some

Euclidean time interval t_1 between the two circles. Unlike for the states given by Eqns. (1)-(3), this path integral is not generally separable into a product of wave functions (4), but, after integrating over time t_1 , it gives the density matrix for a mixed quantum state, and is equal to the propagator

$$K(x_n, 0; x'_n, t_1) = \prod_{n>0, i} \sum_{m_n^{(i)} > 0} \Phi_{m_n^{(i)}}(Y_n^{(i)}) \Phi_{m_n^{(i)}}(Y_n^{(i)'}) e^{-nm_n^{(i)} t_1}, \quad (5)$$

where $\Phi_{m_n^{(i)}}(Y_n^{(i)})$ would match the excited eigenstates of the harmonic oscillators ($\Phi_{m_n^{(i)}} \equiv \Psi_{m_n^{(i)}}$) if we allowed an unlimited resolution for the fields, or equal some wave functions which contain only that part of the information contained in the $\Psi_{m_n^{(i)}}$ that associates with the finite eigenenergy sector surviving after the introduction of a given cut off at a finite highest energy scale. If the handles are off shell, then, instead of the quantum constraint equation (3), we must use the "time-independent" wave equation [8,10]

$$(H_{WDW} - \sum_{n>0, i} nm_n^{(i)}) \Psi = 0, \quad (6)$$

so that the wave function for the wormhole handles becomes

$$\Psi \equiv \Psi(x_0, Y) = e^{iKx_0} \prod_{n>0, i} e^{-\frac{1}{2}n(Y_n^{(i)})^2}. \quad (7)$$

The K^2 term in (6), resulting from the application of operator $-\frac{\partial^2}{\partial x_0^2}$ to Ψ , gives the energy of the x_0 plane waves, with $K^2 < 0$. This would correspond to a timelike momentum in *Lorentzian* space-time. Wick rotating to Euclidean momentum, $K \rightarrow -iK_E$, we have from (6) and (7) $K_E^2 = \sum_{n>0, i} (m_n^{(i)} + \frac{1}{2})n$ and, since we are dealing with harmonic oscillators, we have in general $K_E = \sum_{n>0, i} m_n^{(i)}n$ [11]. Let us then calculate the quantity

$$\begin{aligned} & \sum_{m_n^{(i)} > 0} \Psi(x_0, Y) \Psi^*(x'_0, Y') \\ &= \sum_{m_n^{(i)} > 0} \prod_{n>0, i} (e^{-\frac{1}{2}n(Y_n^{(i)})^2} e^{-\frac{1}{2}n(Y_n^{(i)'})^2} e^{-m_n^{(i)}n(x'_0 - x_0)}). \end{aligned} \quad (8)$$

The relative minus sign between x_0 and x'_0 in (8) should be kept anyway in order to ensure an orientable surface when the handles are glued together [2]. Taking $x'_0 - x_0 = t_1$, noting that, since each two circles can have any time separations, one should integrate (8) over all possible values of t_1 [8], and denoting the density matrix by ρ , we can see that

$$i \int dt_1 K(x_n, 0; x'_n, t_1)$$

$$= \sum_{m_n^{(i)} > 0} \int d(x'_0 - x_0) \Psi(x_0, Y) \Psi^*(x'_0, Y') \equiv i\rho(Y; Y'), \quad (9)$$

whenever we take for the states $\Phi_{m_n^{(i)}}$ in the propagator (5) only that part of the $\Psi_{m_n^{(i)}}$ which corresponds to the harmonic oscillator ground states, surviving after projecting off all the information contained in the Hermite polynomials.

From (7) and (9), the density matrix for handles becomes

$$\rho(Y; Y') = \sum_{m_n^{(i)} > 0} \prod_{n > 0, i} \frac{\Psi_0(Y_n^{(i)}) \Psi_0(Y_n^{(i)'})}{nm_n^{(i)}}, \quad (10)$$

where $\Psi_0(Y_n^{(i)}) = e^{-\frac{1}{2}n(Y_n^{(i)})^2}$. Thus, for each $m_n^{(i)} > 0$, we should use a Green function for the density matrix of the mixed state case given by

$$\begin{aligned} & < x_\mu(z_1).x_\nu(z_2)\dots > \\ &= \int d[x_\mu] x_\mu(z_1) x_\nu(z_2) \dots \prod_r \prod_{n > 0, i} \tilde{\Theta}(x_0 - x'_0) \Psi_0(Y_n^{(i)}) \Psi_0(Y_n^{(i)'}) e^{-I}, \end{aligned} \quad (11)$$

where we have replaced the full δ function in (1) for the density matrix (7), specialising to a single generic relative probability $(nm_n^{(i)})^{-1}$ for each mode $n > 0$, and the step function $i\tilde{\Theta}(x_0 - x'_0) = \Theta(x_0 - x'_0)$ arises from integrating the δ function for the zero mode over its argument, as it is done in (9) and (10). In the present approach, if we want to consider a Green function also for the wave function (7), instead of the full δ function, we should use the probability $|\Psi|^2$ obtained from (7) as the weighting factor. This Green function will then be

$$\begin{aligned} & < x_\mu(z_1).x_\nu(z_2)\dots > \\ &= \int d[x_\mu] x_\mu(z_1) x_\nu(z_2) \dots \prod_r \prod_{n > 0, i} \delta(x_0 - x'_0) \Psi_0(Y_n^{(i)}) \Psi_0(Y_n^{(i)'}) e^{-I}. \end{aligned} \quad (12)$$

If we regard each pair of circles as the ends of a sum of wormholes of different species [1,3], then each species would now be labelled by just the momentum K of the zero mode, but *not* the levels $m_n^{(i)}$ of the other modes. The quantity K can be interpreted as the conserved scalar charge carried by the wormhole [7].

We can now calculate the effect of wormholes on tachyonic amplitudes for handles whose quantum state is given by both a density matrix and a wave function, using the procedure devised by Lyons and Hawking [2]. For the case of handles in mixed state and tachyons with momenta p_j , unlike the pure-state case considered

in Ref. 2, the path integral describing the interaction cannot be factorised into path integrals on each of the two circles [8]. Instead of the wave function (4), one should then introduce as weighting factor the density matrix element $\rho_{m_n}^{(i)}$ which corresponds to each relative probability $\frac{1}{nm_n^{(i)}}$. In the limit of small circle radius $r \rightarrow 0$, we then have for each of these density matrix elements

$$\begin{aligned} D(\rho; p_j) &\propto \int dx_0 dx'_0 \left(\prod_{n>0} dY_n^{(i)} dY_n^{(i)'} \right) \rho_{m_n^{(i)}} \\ &\times \int dr r^{-3+(\sum_1^M p_j)^2} \int [D\zeta_j] \exp[i(x_0 - x'_0) \sum_1^M p_j] \\ &\times \prod_{n>0} e^{\sum_i [-\frac{1}{2}n((Y_n^{(i)})^2 + (Y_n^{(i)'})^2) + ir^n k_n^{(i)}(Y_n^{(i)} - Y_n^{(i)'})]}, \end{aligned} \quad (13)$$

where

$$\rho_{m_n^{(i)}} = \prod_{n>0} \frac{\Psi_0(Y_n^{(i)}) \Psi_0(Y_n^{(i)'})}{nm_n^{(i)}},$$

and we take, as in [2], $r = |z_2|^{-1}$, $\zeta_j = \frac{z_j}{z_2}$, $j = 3, \dots, M$, with M the number of on-shell tachyon vertex operators inserted in the region exterior to the circles, $\zeta_0 = 0$, $\zeta_1 = \infty$, $\zeta_2 = 1$. $\int [D\zeta_j]$ denotes integration over ζ_j with a measure whose explicit form need not be known for our calculation, and $k_n = 2 \sum_{j=1}^M p_j \zeta_j^{-n}$, with $k_n^{(1)}$ and $k_n^{(2)}$ the real and imaginary parts of k_n , respectively.

Each integral pair over $Y_n^{(i)}$ and $Y_n^{(i)'}$ gives a factor $\frac{e^{-\frac{r^{2n}(k_n^{(i)})^2}{n}}}{n}$ for each n . The Gaussian exponential factor would only contribute higher-order interactions and will be disregarded in our calculation [2]. The other possible contribution would come from integration over each pair of zero-mode fields x_0, x'_0 . Since all possible dependence of the density matrix on such field has already been integrated out, unlike for handles in a pure state, we are left with a single delta on $\sum_{j=1}^M p_j$, so in first order approximation the path integral (13) gives essentially a factor $\int \frac{dr}{r^3 m_n^{(i)} n^2}$ for each $m_n^{(i)}$ and n . Note that this factor contains no integration over momentum K . As in the space-time wormhole case [4], the effect of handles will be given again by a vertex operator, located approximately at the center of the circles [1,3]. Thus, for each $m_n^{(i)}$ and n , the stringy wormholes in mixed state will give rise to a bi-local effective action [1] for each $m_n^{(i)}$

$$-\frac{1}{2} \int d\sigma_1 \int d\sigma_2 \sum_{q,i} \frac{V_q(\sigma_1) V_q(\sigma_2)}{m_n^{(i)} n^2}, \quad (14)$$

where V_q are the vertex functions. Following hereafter the same procedure as for wormholes in space-time [4], this action can be

made local by introducing α parameters, i.e.

$$\int d\sigma \sum_{q,i} ((-\frac{1}{2}\alpha_q^2 m_n^{(i)} n^2) + \alpha_q V_q(\sigma)) \quad (15)$$

which leads to a probability measure over the α parameters for each $m_n^{(i)}$

$$Z(\alpha) \prod_{q,i} e^{-\frac{1}{2}\alpha_q^2 m_n^{(i)} n^2}, \quad (16)$$

where again $Z(\alpha)$ is the path integral over all fields x^μ on the two-sphere [1,3], containing the effective interaction $\alpha_q V_q(\sigma)$. The distribution for α -parameters associated with (16) corresponds to just one of the infinite relative probabilities for the state $\Psi_0(Y_n^{(i)})$ of handles. Therefore, one should now sum (16) over all $m_n^{(i)}$ [12,13], to finally obtain a probability measure

$$Z(\alpha) \prod_q (e^{\frac{1}{2}n^2\alpha_q^2} - 1)^{-1} \quad (17)$$

for each n and i . Thus, as it was suggested for 4-dimensional space-time [12,13], we obtain a Planckian distribution for α parameters that allows to interpret $\frac{1}{2}\alpha_q^2$ as the energy of the quanta of a radiation field, and n^{-2} as some temperature, on string-theory superspace.

In the case that the quantum state of the handles be given by the wave function (7), the calculation is similar, but with $\rho_{m_n^{(i)}}$ replaced by $|\Psi(x_0, Y)|^2$ in the path integral (13). In actual calculation, the only difference is in the integration over the field zero modes which now produces $\delta(K - \sum_{j=1}^M p_j)$. The essential factor becomes then $-\frac{1}{n(K^2-2)}$ for each n . It follows that each mode n contributes a probability measure

$$Z(\alpha) \prod_{q,i} e^{-\frac{1}{2}\alpha_q^2(1-\frac{K^2}{2})n} = Z(\alpha) \prod_{q,i} e^{-\frac{1}{2}\alpha_q(K)^2n^2}, \quad (18)$$

where $\alpha_q(K)^2 = \alpha_q^2 \frac{(1-\frac{K^2}{2})}{n}$.

Note that, since the α parameters in both (17) and (18) are labelled by the momentum K , but *not* the levels $m_n^{(i)}$, our results do not allow any interpretation of the α in terms of a quantum field on superspace, which is dimensionally reducible to an infinite tower of fields on space-time. For handles whose state is given by (7), if the initial state is a state with definite values of the α parameters, the final state will be the same as the initial state [14], according to our results, the radiation field α can be dimensionally reduced to just one field, rather than a tower of fields, on the usual

space-time, i.e. on the preferred minisuperspace from string-theory superspace, consisting of just the $n = 0$ modes [1,3]. Therefore, there will always exist a set of classical values for α which makes it possible to drive a consistent mechanism that fixes the values of the coupling constants.

Acknowledgements. This work was supported by CAICYT under Research Project N PB91-0052.

References

- 1** S.W. Hawking, Nucl. Phys. B363, 117 (1991).
- 2** A. Lyons and S.W. Hawking, Phys. Rev. D44, 3802 (1991).
- 3** S.W. Hawking, Phys. Script. T36, 222 (1991).
- 4** S.W. Hawking, Phys. Rev. D37, 904 (1988).
- 5** S.W. Hawking, Nucl. Phys. B335, 155 (1990).
- 6** S. Coleman, Nucl. Phys. B307, 867 (1988).
- 7** A. Lyons, Ph. D. Thesis, University of Cambridge, UK, 1991.
- 8** P.F. González-Díaz, Nucl. Phys. B351, 767 (1991).
- 9** S. Mandelstam, in *Unified Field Theories, Proceedings of the 1985 Santa Barbara Workshop*, eds. M. Green and D. Gross (World Scientific, Singapore, 1986).
- 10** S.W. Hawking, in *300 Years of Gravitation*, eds. S.W. Hawking and W. Israel (Cambridge Univ. Press, Cambridge, 1987).
- 11** W.H. Louisell, *Radiation and Noise in Quantum Electronics* (McGraw Hill, New York, 1964).
- 12** P.F. González-Díaz, Mod. Phys. Lett. A8, 1089 (1993).
- 13** P.F. González-Díaz, Class. Quant. Grav. 10, 2505 (1993).
- 14** S. Coleman, Nucl. Phys. B310, 643 (1988).